

ARL 66-0205  
OCTOBER 1966




## **Aerospace Research Laboratories**

AD 647322  
SIMULTANEOUS TESTS FOR THE EQUALITY OF  
VARIANCES AGAINST CERTAIN ALTERNATIVES

(Reprint from The Australian Journal of Statistics, Vol. 7, No. 3,  
pp. 105-109, November 1965)

P. R. KRISHNAIAH

APPLIED MATHEMATICS RESEARCH LABORATORY



Distribution of this document is unlimited

**OFFICE OF AEROSPACE RESEARCH**  
**United States Air Force**



**ARCHIVE COPY**

## NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Qualified requesters may obtain copies of this report from the Defense Documentation Center, (DDC), Cameron Station, Alexandria, Virginia.

(Reproduction in whole or in part is permitted for any purpose of the U.S. Gov't)

**Distribution of this document is unlimited**

Copies of ARL Technical Documentary Reports should not be returned to Aerospace Research Laboratories unless return is required by security considerations, contractual obligations or notices on a specified document.

**ACCOUNTING BY**

<b>GRSTH</b>	<b>WRITE SECTION /</b>
<b>DGC</b>	<b>DEF SECTION</b>
<b>GRAMMERCED</b>	<input type="checkbox"/>
<b>JUSTIFICATION</b>	<input type="checkbox"/>

**-**

**DISCONTINUATION/INADVERTENT CANCE?**

**BEST.**      **RUAL. end/w GRSTAL**

**✓**      **W**

Reprinted from *The Australian Journal of Statistics*, Vol. 7, No. 3,  
pages 105-109, November, 1965

---

---

**SIMULTANEOUS TESTS FOR THE EQUALITY OF  
VARIANCES AGAINST CERTAIN ALTERNATIVES**

---

**P. R. KRISHNAIAH**

*Aerospace Research Laboratories, Wright-Patterson A.F.B., Ohio*

---

---

## SIMULTANEOUS TESTS FOR THE EQUALITY OF VARIANCES AGAINST CERTAIN ALTERNATIVES<sup>1</sup>

P. R. KRISHNAIAH

*Aerospace Research Laboratories, Wright-Patterson A.F.B., Ohio*

### 1. Introduction

In many situations, the experimenter is interested in testing which of the variances differ significantly from others when the overall hypothesis of equality of variances is rejected. The simultaneous test procedures play an important role in these situations. Hartley (1950) considered the problem of pairwise comparisons of variances simultaneously. Gnanadesikan (1959) considered the problem of comparing several variances simultaneously against a standard one. In the present paper, test procedures are proposed for comparing each variance simultaneously against the next one by using the union-intersection principle of Roy (1953).

### 2. Simultaneous Test Procedures

Consider  $k$  normal populations with means (known or unknown)  $\mu_1, \mu_2, \dots, \mu_k$  and unknown variances  $\sigma_1^2, \sigma_2^2, \dots, \sigma_k^2$ . Let  $s_i^2$  be the sample estimate of  $\sigma_i^2$  based on  $n_i$  degrees of freedom. In addition, let  $H: \sigma_1^2 = \sigma_2^2 = \dots = \sigma_k^2$ . The total hypothesis  $H$  can be expressed as  $H = \bigcap_{i \neq j=1}^k H_{ij}$ ,  $H = \bigcap_{i=1}^k H_{ik}$  and  $H = \bigcap_{i=1}^{k-1} H_{i,i+1}$  where  $H_{rr}: \sigma_r^2 = \sigma_s^2$ . Now, let  $A_1 = \bigcup_{i \neq j=1}^k A_{ij}$ ,  $A_2 = \bigcup_{i=1}^{k-1} A_{ik}$  and  $A_3 = \bigcup_{i=1}^{k-1} A_{i,i+1}$  where  $A_{rr}: \sigma_r^2 \neq \sigma_s^2$ .

Hartley (1950) considered the problems of testing  $H$  against  $A_1$  when the sample sizes are equal. This test is known as the  $F_{\max}$  test. Ramachandran (1956) showed that Hartley's  $F_{\max}$  test is unbiased. The tables which are useful in the application of this test procedure are given in David (1952).

Gnanadesikan (1959) considered the problems of testing  $H$  against  $A_2$ ; his proof for the monotonicity of the power of the test is not correct. We now discuss a one-sided version of the test considered in Gnanadesikan (1959).

<sup>1</sup> Manuscript received June 24, 1965; revised October 25, 1965

A procedure for testing  $H_{1k}, H_{2k}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives  $A_{1k}, A_{2k}, \dots, A_{k-1,k}$  and  $A_2^*$ , where  $A_2^* = \bigcup_{i=1}^{k-1} A_{ik}^*$  and  $A_{ik}^*: \sigma_i^2 > \sigma_k^2$  is discussed here.

According to this procedure, we accept or reject  $H_{ik}$  according as

$$F_{ik} \leq c_{ik}$$

where  $c_{ik}$ 's are chosen such that

$$(2.1) \quad P[F_{ik} \leq c_{ik}; i=1, 2, \dots, k-1 | H] = (1-\alpha), \text{ and} \\ F_{ij} = s_i^2/s_j^2.$$

The total hypothesis  $H$  is accepted if all the individual hypotheses  $H_{1k}, H_{2k}, \dots, H_{k-1,k}$  are accepted. This test procedure is similar to the test proposed by Ghosh (1955) for testing several hypotheses simultaneously under the ANOVA Model I. For practical purposes, we choose the critical values  $c_{ik}$  to be equal, and call them  $c_\alpha$ . When  $n_1 = n_2 = \dots = n_k = n$ , we can obtain the critical values  $c_\alpha$  from the tables of Gupta (1963) for  $\alpha = 0.25, 0.10, 0.05, 0.01, n = 2(2)50$  and  $k = 2(1)11$ . When  $n_1 = n_2 = \dots = n_{k-1} = n$ , we can obtain the critical values  $c_\alpha$  from the tables of Armitage and Krishnaiah (1964) for  $\alpha = 0.10, 0.05, 0.025, 0.01, n = 1(1)19, k = 2(1)13$  and  $n_k = 5(1)45$ . The simultaneous confidence intervals associated with the above test procedure are given by

$$(3.2) \quad P\left[\frac{\sigma_k^2}{\sigma_i^2} \leq \frac{s_k^2}{s_i^2} c_{ik}; i=1, 2, \dots, k-1\right] = (1-\alpha)$$

where  $c_{ik}$ 's are given by (2.1). We will now discuss about another one-sided version of the test considered in Gnanadesikan (1959).

Let  $A_2^{**} = \bigcup A_{ik}^{**}$  where  $A_{ik}^{**}: \sigma_i^2 < \sigma_k^2$ . A procedure is proposed in Krishnaiah and Armitage (1964) to test  $H_{1k}, H_{2k}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives  $A_{1k}^{**}, A_{2k}^{**}, \dots, A_{k-1,k}^{**}$  and  $A_2^{**}$ . When  $n_1 = n_2 = \dots = n_k = n$ , the 25%, 10%, 5% and 1% critical values which are useful in the application of this test procedure can be obtained from the tables of Gupta and Sobel (1962) for  $n = 2(2)50$  and  $k = 2(1)11$ . The 10%, 5%, 2.5% and 1% critical values for this test procedure can be obtained from the tables of Krishnaiah and Armitage (1964) for  $k = 2(1)13, n_k = 5(1)45$  and  $m = 1(1)20$  when  $n_1 = n_2 = \dots = n_{k-1} = m$ . The simultaneous confidence intervals associated with this test procedure are given by

$$(2.3) \quad P\left[\frac{\sigma_k^2}{\sigma_i^2} \geq \frac{s_k^2}{s_i^2} d_{ik}; i=1, 2, \dots, k-1\right] = (1-\alpha)$$

where  $d_{ik}$ 's are chosen such that

$$(2.4) \quad P[F_{ik} \geq d_{ik}; i=1, 2, \dots, k-1 | H] = (1-\alpha).$$

We will now propose a test procedure for testing  $H_{1k}, H_{2k}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives  $A_{1k}, A_{2k}, \dots, A_{k-1,k}$  and  $A_3$ . According to this procedure, we accept  $H_{i,i+1}$  if

$$(2.5) \quad f_{ik} \leq F_{i,i+1} \leq g_{ik}$$

and reject  $H_{i,i+1}$  otherwise. The constants  $f_{ik}$  and  $g_{ik}$  are chosen such that

$$(2.6) \quad P[f_{i\alpha} \leq F_{i,i+1} \leq g_{i\alpha}; i=1, 2, \dots, k-1 | H] = (1-\alpha).$$

The optimum choice of the critical values  $f_{i\alpha}$  and  $g_{i\alpha}$  is not known. For practical purposes, we impose the restriction that the acceptance regions (2.5) are of equal size. In addition, we impose the restriction that, for each  $i$ , the test with the acceptance region (2.5) is locally unbiased.

When  $H$  is true,  $n_1 s_1^2, n_2 s_2^2, \dots, n_k s_k^2$  are distributed independently as central chi-square variates with  $n_1, n_2, \dots, n_k$  degrees of freedom. Starting from the joint distribution of  $s_1^2, s_2^2, \dots, s_k^2$ , making the transformations  $F_{i,i+1} = s_i^2/s_{i+1}^2$ ,  $s_k^2 = s_k^2$  and integrating out  $s_k^2$  we obtain the following expression for the joint frequency function of  $F_{12}, F_{23}, \dots, F_{k-1,k}$  when  $H$  is true:

$$(2.7) \quad f(F_{12}, F_{23}, \dots, F_{k-1,k} | H) = \frac{(n_1/n_k) \left[ \prod_{j=1}^{k-1} n_j \prod_{i=j}^{k-1} F_{i,i+1}/n_k \right]^{1/2} (n_k-2)}{\prod_{j=1}^k \Gamma(n_j/2) [1 + n_k^{-1} \sum_{j=1}^{k-1} n_j \prod_{i=j}^{k-1} F_{i,i+1}]^{1/2} \Sigma n_j}$$

The simultaneous confidence intervals associated with the above test are given by

$$(2.8) \quad P[f_{i\alpha} s_{i+1}^2/s_i^2 \leq \sigma_{i+1}^2/\sigma_i^2 \leq g_{i\alpha} s_{i+1}^2/s_i^2; i=1, 2, \dots, k-1] = (1-\alpha).$$

The above simultaneous confidence intervals can be derived by using the fact that

$$(2.9) \quad P[f_{i\alpha} \leq F_{i,i+1} \leq g_{i\alpha}; i=1, 2, \dots, k-1 | H] = P[f_{i\alpha} \leq \sigma_{i+1}^2 F_{i,i+1} / \sigma_i^2 \leq g_{i\alpha}; i=1, 2, \dots, k-1 | \cap A_i] = (1-\alpha).$$

A procedure is proposed below for testing  $H_{12}, H_{23}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives

$$A_{12}^*, A_{23}^*, \dots, A_{k-1,k}^* \text{ and } A_3^* \text{ where } A_3^* = \bigcup_{i=1}^{k-1} A_{i,i+1}^* \text{ and } A_{i,i+1}^*: \sigma_i^2 > \sigma_{i+1}^2$$

According to this procedure, we accept or reject  $H_{i,i+1}$  according as

$$F_{i,i+1} \leq a_{i\alpha}$$

where  $a_{i\alpha}$ 's are chosen such that

$$(2.10) \quad P[F_{i,i+1} \leq a_{i\alpha}; i=1, 2, \dots, k-1 | H] = (1-\alpha).$$

The simultaneous confidence intervals associated with the above test are given by

$$(2.11) \quad P[\sigma_{i+1}^2/\sigma_i^2 \leq a_{i\alpha} s_{i+1}^2/s_i^2; i=1, 2, \dots, k-1] = (1-\alpha).$$

For practical purposes, we choose the critical values  $a_{i\alpha}$  in (2.10) to be equal.

We test  $H_{12}, H_{23}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives  $A_{12}^{**}, A_{23}^{**}, \dots, A_{k-1,k}^{**}$  and  $A_3^{**}$  where

$$A_3^{**} = \bigcup_{i=1}^{k-1} A_{i,i+1}^{**} \text{ and } A_{i,i+1}^{**}: \sigma_i^2 < \sigma_{i+1}^2, \text{ by accepting or rejecting } H_{i,i+1} \text{ according as}$$

$$F_{i,i+1} > b_{i\alpha}$$

where  $b_{i\alpha}$ 's are chosen such that

$$(2.12) \quad P[F_{i,i+1} \geq b_{i\alpha}; i=1, 2, \dots, k-1 | H] = (1-\alpha).$$

The simultaneous confidence intervals associated with the above test are given by

$$(2.13) \quad P\left[\frac{\sigma_{i+1}^2}{\sigma_i^2} \geq \frac{s_{i+1}^2}{s_i^2} b_{i\alpha}; i=1, 2, \dots, k-1\right] = (1-\alpha).$$

As before we can, for practical purposes, choose the critical values  $b_{i\alpha}$ 's to be equal. When  $k=3$ , the values of  $\alpha$  in (2.12) can be computed for given values of  $b_{i\alpha}$  and  $n_i$  by using the method discussed in Bechhofer and Sobel (1954) and Rozivich, Bancroft and Hartley (1956).

### 3. General Remarks

Hartley's  $F_{max}$  is useful when the experimenter is interested in testing the hypotheses  $H_{rs}$  ( $r \neq s=1, 2, \dots, k$ ) simultaneously and the sample sizes are equal. But there are many situations when the experimenter is interested in testing a subset of these hypotheses. In these situations, it is not desirable to use the  $F_{max}$  test since alternative simultaneous test procedures yield shorter (in terms of lengths) confidence intervals. For example, if we are interested in testing  $H_{12}, H_{23}, \dots, H_{k-1,k}$  and  $H$  simultaneously against the respective alternatives  $A_{12}, A_{23}, \dots, A_{k-1,k}$  and  $A_3$ , the lengths of the confidence intervals associated with the simultaneous test procedure proposed in this paper are shorter than those associated with the  $F_{max}$  test if the critical values  $f_{i\alpha}$  and  $g_{i\alpha}$  in (2.6) are chosen such that  $g_{i\alpha} = g_\alpha$  and  $f_{i\alpha} = g_\alpha^{-1}$  for  $i=1, 2, \dots, k-1$  when the sample sizes are equal. Similarly, if we choose the critical values properly, the lengths of the confidence intervals associated with Gnanadesikan's test are shorter than the lengths of the corresponding confidence intervals associated with the  $F_{max}$  test.

Now let  $\beta$  denote the Type II error associated with the test proposed in this paper for testing  $H$  against  $A_2^*$ . When  $\bigcap_{i=1}^{k-1} A_{ik}^*$  is true,

$$\begin{aligned} 1-\beta &= P[F_{ik} \geq c_{i\alpha}; i=1, 2, \dots, k-1 | \bigcap_{i=1}^{k-1} A_{ik}^*] \\ &= \int_{c_{1,\alpha}/\delta_{1,k}}^{\infty} \dots \int_{c_{k-1,\alpha}/\delta_{k-1,k}}^{\infty} f(F_{1,k}, \dots, F_{k-1,k} | H) \prod_{i=1}^{k-1} dF_{i,k} \end{aligned}$$

where  $\delta_{i,k} = \sigma_i^2/\sigma_k^2$ . It is now evident that  $(1-\beta)$  increases as each non-centrality parameter  $\delta_{i,k}$  increases. The above proof can be easily modified when only some of the  $A_{ik}^*$ 's are true. We can similarly show that other one-sided test procedures considered in this paper are monotonic increasing functions of the non-centrality parameters.

The test procedures discussed in the present paper are analogous to some special cases of the tests considered in Krishnalah (1965a, 1965b) for multiple comparisons of means.

## References

- Armitage, J. V., and Krishnaiah, P. R. (1964). *Tables for the Studentized Largest Chi-Square Distribution and Their Applications*. ARL 64-188, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio.
- Bechhofer, R. E., and Sobel, M. (1954). "A single sample multiple decision procedure for ranking variances of normal populations." *Ann. Math. Statist.*, 25, 275-286.
- Bozovich, H., Bancroft, T. A., and Hartley, H. O. (1956). "Power of analysis of variance test procedures for certain incompletely specified models, I." *Ann. Math. Statist.*, 27, 1017-1043.
- David, H. A. (1952). "Upper 5 and 1% points of the maximum F-ratio." *Biometrika*, 39, 422-424.
- Ghosh, M. N. (1955). "Simultaneous tests of linear hypotheses." *Biometrika*, 42, 441-449.
- Gnanadesikan, R. (1959). "Equality of more than two variances and more than two dispersion matrices against certain alternatives." *Ann. Math. Statist.*, 30, 177-184.
- Gupta, S. S., and Sobel, M. (1962). "On the smallest of several correlated F statistics." *Biometrika*, 49, 509-523.
- Gupta, S. S. (1963). "On a selection and ranking procedure for gamma populations." *Ann. Inst. Statist. Math.*, 14, 199-216.
- Hartley, H. O. (1950). "The maximum F-ratio as short-cut test for heterogeneity of variances." *Biometrika*, 37, 308-312.
- Krishnaiah, P. R., and Armitage, J. V. (1964). *Distribution of the Studentized Smallest Chi-Square, with Tables and Applications*. ARL 64-218, Aerospace Research Laboratories, Wright-Patterson Air Force Base, Ohio.
- Krishnaiah, P. R. (1965a). "On the simultaneous ANOVA and MANOVA tests." *Ann. Inst. Statist. Math.*, 17, 50-53.
- Krishnaiah, P. R. (1965b). "Multiple comparison tests in multi-response experiments." *Sankhyā*, 27.
- Ramachandran, K. V. (1956). "On the Tukey test for the equality of means and the Hartley test for the equality of variances." *Ann. Math. Statist.*, 27, 825-831.
- Ramachandran, K. V. (1956). "Contributions to simultaneous confidence interval estimation." *Biometrics*, 12, 51-56.
- Roy, S. N. (1953). "On a heuristic method of test construction and its uses in multivariate analysis." *Ann. Math. Statist.*, 24, 220-238.